

Performance of eight tree species in the highland Vertisols of central Ethiopia: growth, foliage nutrient concentration and effect on soil chemical properties

KINDU MEKONNEN^{1,*}, TADESSE YOHANNES²,
GERHARD GLATZEL¹ and YOSEF AMHA²

¹Institute of Forest Ecology, University of Natural Resources and Applied Life Sciences, Peter-Jordan-Strasse 82, A-1190, Vienna, Austria; ²Ethiopian Agricultural Research Organization (EARO), P.O. Box 2003 Holetta, Ethiopia; *Author for correspondence (e-mail: kindumeko@yahoo.com; phone: +43-1-47654-4117; fax: +43-1-4797896)

Received 6 July 2004; accepted in revised form 7 February 2006

Key words: Available P, Biomass, Foliage to stem ratio, Foliage and soil N, Height

Abstract. The coverage of trees in the highland Vertisol areas of Ethiopia is very scarce. A tree screening trial was conducted from 1997 to 2002 in Ginchi (central Ethiopia) to select fast growing and high biomass producing tree species; evaluate foliage macronutrient concentration of different tree species; and assess effects of trees on soil chemical properties beneath their canopies. *Acacia decurrens* Willd., *A. mearnsii* De Wild and *Eucalyptus globulus* Labill attained the highest height growth at 64 months as compared to other indigenous and exotic species. *E. globulus* provided better height increment from 24 to 36 and 36 to 64 months than other tree and shrub species. *Acacia mearnsii* and *A. saligna* Labill Wendl produced high biomass at 40 and 64 months. Differences between the highest and lowest dry biomass at 12, 40 and 64 months were 1.13, 29.19 and 38.89 kg tree⁻¹, respectively. None of the tree species resulted in a foliage to stem biomass ratio of >0.98 at 40 and 64 months. *Sesbania sesban* (L.) Merr had high N and P concentrations in its foliages and stems at 12 and 40 months. Total N under *Acacia abyssinica* Hochst. Ex Benth, *A. saligna* and *S. sesban* was slightly greater at 40 months than 12 months.

Introduction

Vertisols are the fourth most important soil orders and cover over 10% (12.7 million ha) of the Ethiopian landmass (Berhanu 1985). From the total coverage of Vertisols, 7.6 million ha is found in the highlands (> 1500 masl). The distribution and diversity of trees and shrubs on Vertisols is very scarce due to mainly anthropogenic, climatic and edaphic factors (EARO (Ethiopian Agricultural Research Organization) 2000). Farmers establish *E. globulus* woodlot around the homesteads and use it for fuelwood, poles and as sources of cash. Other naturally grown *Acacia* species exist in riverbanks, grazing and croplands with a very little area coverage.

Waterlogging in the rainy season and cracking in the dry season are features of Vertisols. The cracks on Vertisols are deep and wide. Preliminary observation at Ginchi (central Ethiopia) showed up to 1.06 m depth and 0.12 m width of soil cracks on fallow lands and 0.67 m depth and 0.09 m width of

cracks on cultivated lands (Worku 1997, personal communication). When the soil form cracks tree and shrub roots break and cease anchorage and water absorption activities. This has resulted in poor adaptation of trees and shrubs on Vertisols.

Shortages of wood and improved animal feed, soil erosion, decline of soil fertility and lack of alternative fuel wood sources are critical problems in the highland Vertisol areas (ICRAF (International Center for Agroforestry Research) 1990). Farmers on Vertisol areas demand wood mainly for fuel wood, construction poles, farm implements and charcoal making. In 1992 total requirements for wood products in the country was estimated to be 47.4 million m³ (EFAP (Ethiopian Forestry Action Program) 1994). In 2014 total wood requirement and supply with the implementation of programs developed by EFAP is estimated to be 30.2 and 94.6 million m³, respectively. Wood requirement outweighs the supply both in Vertisol and other areas of the country.

Soil erosion and continuous cultivation with no or minimum inputs have caused depletion of soil nutrients on Vertisols. Hague et al. (1993) reported deficiencies of N and P on highland Vertisols. Correcting N and P deficiencies by fertilizer application alone is a challenging task in view of the high fertilizer cost and low income of the subsistent farmers. Organic resources like crop residues and cow dung are advertised as one of the options for replenishment of soil fertility. However, the attempt is challenged by competing uses of crop residues and manures for animal feed and fuel sources, respectively.

Introduction and improved management of adaptable, multipurpose trees and shrubs can be one of the strategies to minimize the current wood, feed and soil related constraints. So far, available information on the performance of agroforestry tree and shrub species on Vertisols is limited. The objectives of the study were therefore to (i) select fast growing and high biomass producing tree species, (ii) evaluate foliage macronutrient concentration of different tree species and (iii) assess effects of trees on soil chemical properties beneath their canopies.

Materials and methods

Site characteristics

The study was conducted from 1997 to 2002 at Ginchi research sub-center of western Shewa, Ethiopia. The sub-center is geographically located at 09°02' N latitude and 38°07' E longitudes. Altitude is 2200 masl. The sub-center receives rain in the short and long cropping seasons. Mean annual rainfall is 1042 mm. The annual mean maximum and minimum temperatures are 25 and 7 °C, respectively. The soil is classified as Pellic Vertisol. Some of the chemical and physical properties of the top 0–15 and 15–30 cm soil layers were: pH (1:1 soil/water suspension) = 5.99, organic C (g kg⁻¹) = 22.6, total N (g kg⁻¹) = 1.1,

available P (mg kg^{-1})=6.2, exchangeable Ca (cmol kg^{-1})=37.50, Mg (cmol kg^{-1})=10.50, K (cmol kg^{-1})=1.54 and Clay (%)=76. At 15–30 cm depth, pH=6.2, organic C=20.1, total N=1.00, available P=6.5, exchangeable Ca=39.69, Mg=13.08, K=1.62 and Clay=75.

Seed sources and raising of seedlings

Seeds of different agroforestry trees were obtained from Ethiopian Forestry Research Center (FRC), International Livestock Research Institute (ILRI) and Kenya Forestry Research Institute (KEFRI). Seed treatment techniques such as scarification, nipping, soaking in hot and cold water were exercised depending upon the need for the species. Treated seeds of trees and shrubs were directly sown in a polythene tube that contained a mixture of local topsoil, forest soil and sand. The proportion of the soil mixture was 4 (local soil): 2 (forest soil): 1 (sand). The size of the polythene tube was 15 cm height and 8 cm diameter. All pots sown with respective species were exposed to similar watering, shading, weeding and hardening practices.

Experimental design and management

Treatments (different tree species) laid out in a randomized complete block design (RCBD) with three replications (Gomez and Gomez 1984; Wood and Burley 1991). The tree species were: (i) *Acacia abyssinica* Hochst. Ex Benth (collection from Tateke, western Shewa, Ethiopia); (ii) *A. decurrens* Willd (collection from Sebeta, south-west Shewa, Ethiopia); (iii) *A. melanoxylon* R. Br. (collection from Tateke, western Shewa, Ethiopia); (iv) *A. mearnsii* De Wild (collection from mortu Molosi, Kenya); (v) *A. saligna* Labill Wendl. (collection from Zwai with accession no. 7178, east Shewa, Ethiopia); (vi) *Cordia africana* Lam (collection from Sokoru, Ethiopia); (vii) *Eucalyptus globulus* Labill (collection from Deksis, Ethiopia) and (viii) *Sesbania sesban* (L.) Merr (collection from Zwai with accession no. 10865, eastern Shewa, Ethiopia).

Field preparation began one week before planting with hole digging at depths of 30 cm. Trees were planted on hand made broad bed and furrow (BBF) for the purpose of draining excess water. A plot consisted of five BBFs (Figure 1). The size of a plot was 6.5×6 m. Each BBF was 1.2 m wide from the centers of two neighboring furrows, giving a bed 0.8 m wide for establishment of the trees. The length of each BBF was 6.5 m. The height of the BBF was 0.2 m from the furrow center and each furrow was 0.4 m wide (Tedla et al. 1999). Each BBF had a line of five trees. The trees were planted in the center of the BBF. Distance between trees in the same line was 1.3 m while distance between the tree lines in the same plot was 1.2 m. Distance between plots was 2 m.




Figure 1. Sketch of a plot with five broad bed and furrow (BBF).

Measurement, sampling, and soil and foliage analysis

Initial height of the different species was determined at planting (1997). Similarly, height data was taken 12, 24, 36 and 64 months after planting as described by Wood and Burley (1991). Graduated stick and clinometers were used to measure tree and shrub heights. Systematic sampling was followed to select tree and shrub species for biomass determination. Out of the total 25 trees plot⁻¹, 16 of them were border trees. Destructive tree sampling was done at 12, 40, and 64 months from the nine central trees that were not adjacent each other. In every destructive sampling scheme one tree per plot was cut at the ground level. Harvested trees were separated into foliage (leaves and twigs) and stems. Each part was weighted, chopped and sub-sampled for dry matter determination. Sub-samples were oven dried at 70 °C, ground to pass through 1 mm sieve and analyzed for total N using Kjeldahl procedure (Buresh et al. 1982), P (Murphy and Riley 1962) and extractable K (Richards 1954).

Soil auguring was carried out 12 months after planting at each plot in all the replications. To monitor changes in soil characteristics, additional samples were taken 36 months after planting. Sampling layers in all sampling schemes were 0–0.15, 0.15–0.3, 0.3–0.5, 0.5–1, 1–1.5 and 1.5–2 m depths (Kindu et al. 1997). The samples were analyzed for total N following the description by Keeney and Nelson (1982). Available P was determined by Olsen method (Olsen and Sommers 1982).

Statistical analysis

The following formula was used to calculate dry weight of the aboveground tree components.

$$\text{Dry weight}_{\text{total}} = \left[\frac{\text{Fresh weight}_{\text{total}} * \text{Dry weight}_{\text{sample}}}{\text{Fresh weight}_{\text{sample}}} \right] \quad (1)$$

Foliage to stem ratio was calculated based on the relationship between dry foliage and stem biomass. Mean values were used for plotting height, height increment, total soil N and available soil P graphs. The general linear model procedure of the SAS Program (SAS Institute 1999–2001) was used for data analysis. Means for height growth, height increment, aboveground dry biomass and foliage nutrient (N, P and K) concentrations of the different trees were compared using the least significance difference (LSD) at $p < 0.01$.

Results

Growth of trees and shrubs

The height of *A. mearnsii* was higher at 12 months ($p < 0.01$) than the other tree and shrub species (Table 1 and Figure 2a). On the other hand, *A. abyssinica* demonstrated the lowest height growth at 12 months. *A. decurrens* and *A. mearnsii* provided better height growth at 24, 36 and 64 months. *Eucalyptus globulus*, *A. decurrens* and *A. mearnsii* attained the highest growth at 64 months as compared to other indigenous and exotic tree and shrub species. Indigenous trees such as *A. abyssinica* and *C. africana* showed slow growth rates at 12, 24, 36 and 64 months. The growth rate of *S. sesban* was slow after 12 months.

Mean annual height increment of *A. abyssinica* from 12 to 24 and 24 to 36 months was uniform (Table 1 and Figure 2b). *A. decurrens* showed significant height increment from 12 to 24 months. Likewise, *E. globulus* resulted better height increment from 24 to 36 and 36 to 64 months than other tree and shrub species.

Table 1. Comparison of mean height growth and height increment (m) for eight tree species evaluated on a Vertisol area of Ginchi, central Ethiopia.

Tree and shrub species	Height growth				Height increment			
	12 Months	24 Months	36 Months	64 Months	12 vs. 24 months	24 vs. 36 months	36 vs. 64 months	
<i>Acacia abyssinica</i>	0.97 ^c	1.46 ^d	1.97 ^c	4.86 ^{cd}	0.50 ^{dc}	0.50 ^c	2.89 ^{ba}	
<i>Acacia decurrens</i>	2.83 ^{ba}	5.13 ^a	6.89 ^a	9.73 ^a	2.30 ^{at}	1.76 ^{ba}	2.84 ^{ba}	
<i>Acacia mearnsii</i>	3.03 ^a	4.89 ^a	6.64 ^a	9.09 ^{at}	1.86 ^{ba}	1.74 ^{ba}	2.45 ^{bac}	
<i>Acacia melanoxylon</i>	1.76 ^d	2.65 ^c	3.90 ^c	5.65 ^{cb}	0.89 ^c	1.25 ^{bc}	1.75 ^{bc}	
<i>Acacia saligna</i>	2.32 ^c	3.87 ^b	4.99 ^{bc}	6.63 ^b	1.55 ^b	1.12 ^{bc}	1.64 ^{bc}	
<i>Eucalyptus globulus</i>	1.88 ^d	3.59 ^b	6.06 ^{ba}	9.85 ^a	1.71 ^b	2.47 ^a	3.79 ^a	
<i>Cordia africana</i>	1.77 ^d	1.86 ^d	2.39 ^{de}	3.42 ^d	0.09 ^d	0.54 ^c	1.02 ^{bc}	
<i>Sesbania sesban</i>	2.56 ^{bc}	3.49 ^b	3.73 ^{dc}	4.43 ^{cd}	0.94 ^c	0.24 ^c	0.70 ^c	
<i>p</i>	0.0001	0.0001	0.0001	0.0001	0.0001	0.0115	0.0492	

p = significance level of *F*-test, testing the null hypothesis of no difference.

Means in a column followed by the same letter do not differ significantly at their respective *p* value.




Figure 2. Tree species evaluated on a Vertisol area of Ginchi, central Ethiopia (a) mean height and (b) mean height increment.

Biomass production

Dry biomass of the trees widely varied among species. *S. sesban* (L.) Merr provided the highest biomass yield at 12 months. Similarly, *A. mearnsii* and *A. saligna* produced high biomass at 40 and 64 months (Table 2). Differences between the highest and lowest dry biomass at 12, 40 and 64 months were 1.13, 29.19 and 38.89 kg tree⁻¹, respectively. The order for biomass increment of the top four species from 12 to 40 months was *A. decurrens* (29.9) > *A. saligna* (26.8) > *A. mearnsii* (23.7) > *E. globulus* (10.4 kg tree⁻¹). Biomass increment from 40 to 64 months was not as high as that of 12–40 months. The highest biomass increment from 40 to 64 months was 19.26 kg tree⁻¹ for *A. mearnsii* and the lowest was 3.3 kg tree⁻¹ for *C. africana*. *Acacia saligna* and *E. globulus* had a foliage to stem biomass ratio > 1 at 12 months as compared to other species. The highest foliage to stem biomass ratio was 1.18 at 12 months and the lowest was 0.12 at 40 months. None of the tree species resulted in a foliage to stem biomass ratio > 0.98 at 40 and 64 months. The standard deviation (SD) for the total biomass of *A. saligna* and *A. abyssinica* at 64 months was higher than the SD of other tree species.

Foliage nutrient concentration

Foliage N concentration of *S. sesban* was not significantly different ($p < 0.01$) as compared to the foliage N concentration of *A. mearnsii* and *A. melanoxylon* at 12 months (Table 3). The three leguminous tree species (*A. abyssinica*, *A. decurrens* and *A. saligna*) and a none-leguminous species (*C. africana*) did not show significant difference for N concentration in their foliages at 12 months. Foliage N concentration was higher for all species at 40 months than 12 months. Foliage P concentration of *S. sesban* was significantly higher ($p < 0.01$) at 12 and 40 months than most other species. Potassium concentration of foliages of all trees was higher at 40 months than 12 months. *C. africana* followed by *S. sesban* had higher concentration of K both at 12 and 40 months.

Total N and available P trends

The trend of total soil N appeared to be high in the top 0–30 cm depths under all tree species (Figure 3). The stock of total N under *E. globulus* was reduced at 40 months as compared to the stock at 12 months. Total N under *A. abyssinica*, *A. saligna* and *S. sesban* was slightly greater at 40 months than 12 months. Contrary to the N content, the available P content under most species and nearly all soil depths was lower at 40 months than 12 months (Figure 4). Under *A. saligna*, available P below 160 cm was higher at

Table 2. Aboveground dry-biomass of eight tree species evaluated at different months on a Vertisol area of Ginchi, central Ethiopia.

Tree and shrub species	12 Months			40 Months			64 Months		
	Total dry-biomass (kg tree ⁻¹)	SD*	Foliage to stem ratio	Total dry-biomass (kg tree ⁻¹)	SD*	Foliage to stem ratio	Total dry-biomass (kg tree ⁻¹)	SD*	Foliage to stem ratio
<i>Acacia abyssinica</i>	0.21 ^d	0.11	0.50	2.71 ^c	0.90	0.56	13.67 ^{de}	12.22	0.60
<i>Acacia decurrens</i>	1.10 ^{ba}	0.21	0.92	31.02 ^a	6.08	0.36	35.75 ^{ba}	5.20	0.36
<i>Acacia mearnsii</i>	1.03 ^{ba}	0.34	0.75	24.79 ^a	8.77	0.35	44.05 ^a	5.22	0.46
<i>Acacia melanoxylon</i>	0.33 ^d	0.02	0.83	3.90 ^{cb}	2.40	0.98	12.60 ^{de}	1.05	0.87
<i>Acacia saligna</i>	0.80 ^{bc}	0.18	1.05	27.60 ^a	5.37	0.33	43.79 ^a	20.96	0.42
<i>Eucalyptus globulus</i>	0.79 ^{bc}	0.12	1.18	11.18 ^b	4.86	0.46	23.26 ^{bc}	3.34	0.45
<i>Cordia africana</i>	0.56 ^{de}	0.22	0.54	1.83 ^c	0.77	0.12	5.16 ^d	2.84	0.25
<i>Sesbania sesban</i>	1.34 ^a	0.44	0.25	7.83 ^{cb}	3.17	0.20	13.40 ^{de}	5.20	0.31
<i>p</i>	0.0011			0.0001			0.0008		

*Standard Deviation.

p = significance level of *F*-test, testing the null hypothesis of no difference.Means in a column followed by the same letter do not differ significantly at their respective *p* value.

Table 3. Foliage macronutrient content of eight tree species evaluated at different months on a Vertisol area of Ginchi, central Ethiopia.

Tree and shrub species	Nitrogen (g kg ⁻¹)		Phosphorus (g kg ⁻¹)		Potassium (g kg ⁻¹)	
	12 months	40 months	12 months	40 months	12 months	40 months
<i>Acacia abyssinica</i>	22.3 ^{ba}	31.6 ^b	2.33 ^a	2.37 ^b	6.19 ^{cb}	12.50 ^c
<i>Acacia decurrens</i>	20.9 ^{ba}	26.7 ^{cb}	1.23 ^b	1.06 ^c	3.19 ^{cd}	6.67 ^e
<i>Acacia meurnsii</i>	23.7 ^a	28.5 ^b	1.23 ^b	1.17 ^c	5.83 ^{cbd}	7.57 ^{ed}
<i>Acacia melanoxlon</i>	23.6 ^a	24.5 ^{cd}	1.40 ^b	1.25 ^c	6.21 ^{cb}	9.63 ^d
<i>Acacia saligna</i>	20.3 ^{ba}	28.7 ^b	1.43 ^b	1.57 ^c	2.72 ^d	9.10 ^{ed}
<i>Eucalyptus globulus</i>	11.8 ^b	15.9 ^c	1.27 ^b	1.27 ^c	4.08 ^{cd}	9.47 ^d
<i>Cordia africana</i>	18.0 ^{ba}	18.1 ^{ed}	2.57 ^a	2.23 ^b	15.67 ^a	26.03 ^a
<i>Sesbania sesban</i>	28.1 ^a	38.2 ^a	2.80 ^a	3.40 ^a	8.33 ^b	21.23 ^b
<i>p</i>	0.2230	0.0001	0.0010	0.0001	0.0001	0.0001

p = significance level of *F*-test, testing the null hypothesis of no difference. Means in a column followed by the same letter do not differ significantly at their respective *p* value.




Figure 3. Trends of total N at 12 and 40 months under different trees on a Vertisol area of Ginchi, central Ethiopia.

40 months than 12 months. *Acacia mearnsii* and *E. globulus* did not differently impact available P content at 2 m soil depth.

Discussion

Growth and productivity of trees depend upon the genetic, climatic, edaphic and management factors (Lugo et al. 1988). The slow growth performance of the indigenous trees like *A. abyssinica* and *C. africana* can be attributed to the genetic makeup of the species. The annual height and biomass increment of *A. abyssinica* and *C. africana* was as low as other Ethiopian indigenous tree species. Unlike *E. globulus*, *S. sesban* had low height increment from 36 vs. 64 months (Table 1 and Figure 2b). On the other hand, *S. sesban* produced the highest biomass at 12 months as compared to other indigenous and exotic tree and shrub species (Table 2). The continuity of growing and yielding high biomass depends on the lifespan of species. Short lifespan shrub species such as *S. sesban* showed higher annual height and biomass increment in its early than




Figure 4. Trends of available P at 12 and 40 months under different trees on a Vertisol area of Ginchi, central Ethiopia.

late ages. Our findings on height growth and aboveground biomass of *S. sesban* are in line with the findings of Mengistu et al. (2002).

The foliage to stem biomass ratio of many of the species did not increase with time. It has rather decreased as the proportion of the stem dry biomass increased for many of the species. For *E. globulus*, foliage to stem ratio declined from 1.18 in the 12 months to 0.45 in the 64 months (Table 2). During the experimental period, *C. african* shed leaves more than other tree species. The low foliage to stem biomass ratio for *C. african* can be partly explained to leaf shedding. Other tree species such as *A. decurrens* and *A. mearnsii* had feathery and small size leaves. The small size and weight of leaves of the two species could be some of the reasons for the low foliage to stem biomass ratios.

The standard deviation for total dry biomass of *A. saligna* and *A. abyssinica* at the 64 months was 20.9 and 12.2, respectively (Table 2). Increased standard deviation for the two species could be a result of lack of sampling representative trees. Systematic sampling was followed in each experimental plot to select tree and shrub species for biomass determination. We used systematic sampling methods to avoid cutting of adjacently growing trees. Considerations of various criteria reduce extremes from the mean dry biomass of the trees but may lead to cutting of adjacently growing trees. Representative trees for biomass determination can be defined based on diameter, basal area and height distribution of existing stands when the experimental plots consist of more trees than the present study.

The nutrient concentration of tree foliages varies depending on species, soil type and age (Jama et al. 2000). Similarly, the concentration and availability of nutrients under trees vary with depth of soil and distance from a tree base (Mazzarino et al. 1991). Increasing trends of foliage and soil N for *Acacia* species and *S. sesban* in comparison with *E. globulus* can be accounted for the enhancement of biological N fixation of the former than the later. The amount of N fixed by legume tree species can be quite considerable. Ndoye and Dreyfus (1988) estimated the amount of N ($43\text{--}102\text{ kg N ha}^{-1}\text{ y}^{-1}$) that can be fixed by *S. sesban*. Nitrogen concentration of the N fixing tree species was greater than the critical level of $20\text{--}25\text{ g kg}^{-1}$ (Palm et al. 1997). Net immobilization of N can be expected if the species with N concentration below the critical levels are used as organic fertilizer resources.

Soil N under *E. globulus* was less at 40 months than at 12 months. Most *Eucalyptus* species are known for utilizing high amount of resources to fulfill the demand for their fast growth rates (Malik and Sharma 1990). Poor decomposability (higher lignin and soluble polyphenols) of *Eucalyptus* litter compared to litter from other tree species could be the other reason for decline of soil N under *E. globulus*. Likewise, the lowering of soil N in the top 30 cm soil layers under *A. mearnsii* (Figure 3) occurred probably due to less leaf litter deposition of the trees and poor undergrowth of grasses and other annual plants. The absence of leaf litter and poor vegetation undergrowth could have facilitated topsoil erosion and hence N depletion.

Most tree species had low foliage P concentration in comparison with N concentration. The foliage P concentration in *S. sesban* was exceptionally higher at 40 months than the other species. Like the foliage P, soil P under most species declined at 40 months as compared with 12 months. The decline of P in the foliage and the soil can be explained to the very low concentrations of available P in the sub-soil and low root length densities of the species (IAEA (International Atomic Energy Agency) 1975). The addition of available P below 150 cm soil depths under *A. saligna* at 40 months (Figure 4) could be attributable to the association of the tree roots with mycorrhizas. The extensive proliferation of mycorrhizal hyphae enables the exploration of greater soil volume (Lajtha and Harrison 1995).

Conclusions

Based on evaluation of eight tree species on Vertisols, the following were observed: (i) *Acacia mearnsii* and *A. decurrens* provided better height growth at 12, 24, 36 and 64 months, (ii) *Eucalyptus globulus* had shown higher height increment from 24 to 36 and 36 to 64 months than other species, (iii) *S. sesban* had high N and P concentrations in its foliages and stems at 12 and 40 months, (iv) *Acacia mearnsii* and *A. saligna* were superior tree species on Vertisols in terms of biomass production at 40 and 64 months, (v) *A. abyssinica*, *A. saligna* and *S. sesban* slightly improved total N at 40 months as compared to 12 months, (vi) *Acacia mearnsii* lowered soil N in the top 30 cm soil layers and (vii) Available P content under most species and nearly all soil depths was lower at 40 months than 12 months.

Acknowledgements

The authors thank Ethiopian Agricultural Research Organization (EARO) for funding the research work. The support from staff of forestry, soil and water management research divisions at Holetta, and the encouragement from Dr. Demel Teketay are highly appreciated. We would also like to thank anonymous reviewers for their constructive comments.

References

- Berhanu D. 1985. The Vertisols of Ethiopia: their properties, classification and management. In: Fifth meeting of the eastern African sub-committee for soil correlation and land evaluation. World soil resources report no. 56. FAO (Food and Agriculture Organization), Rome. pp. 31–54.
- Buresh R.J., Austin E.R. and Craswell E.T. 1982. Analytical methods in N¹⁵ research. Fert. Res. 3: 37–62.
- EARO (Ethiopian Agricultural Research Organization). 2000. Forestry research strategy. EARO, Addis Abeba, Ethiopia. 107 p.

- EFAP (Ethiopian Forestry Action Program). 1994. The Challenge for Development. Volume II, Addis Abeba, Ethiopia.
- Gomez K.A., Gomez A.A. 1984. Statistical Procedures for Agricultural Research, 2nd ed. IRRA, the Philippines and Wiley, New York, pp. 39–75.
- Haque I., Abebe M., Mamo T. and Dibabe A. 1993. Nutrient management. In: Mamo T., Astatke A., Srivastava K.L. and Dibabe A. (eds), Improved Management of Vertisols for Sustainable Crop-livestock Production in the Ethiopian Highlands: Synthesis Report 1986–92, Technical committee of the joint Vertisol project, Addis Ababa, Ethiopia, pp. 51–73.
- IAEA (International Atomic Energy Agency). 1975. Root activity patterns of some tree crops. Tech. Rep. Ser. No. 170. Int. Atomic Energy Agency, Vienna, Austria.
- ICRAF (International center for Research in Agroforestry). 1990. Agroforestry: Potentials and research needs for the Ethiopian highlands, AFRENA report no. 21, Nairobi, Kenya.
- Jama B., Palm C.A., Buresh R.J., Niang A., Gachengo K., Nziguherba G. and Amadalo B. 2000. *Tithonia diversifolia* as a green manure for soil fertility improvement in western Kenya: a review. *Agroforest. Syst.* 49: 201–221.
- Keeney D.R. and Nelson D.W. 1982. Nitrogen-inorganic forms. In: Page A.L. (ed.), *Methods of Soil Analysis. Agron. 9, Part 2: Chemical and Microbiological Properties*, 2nd ed., Am. Soc. Agron., Madison Wisc., USA, pp. 643–698.
- Kindu M., Buresh R.J. and Jama B. 1997. Root and inorganic nitrogen distributions in *Sesbania* fallow, natural fallow and maize fields. *Plant Soil* 188: 319–327.
- Lajtha K. and Harrision A.F. 1995. Strategies of phosphorus acquisition and conservation by plant species and communities. In: Tiessen H. (ed.), *Phosphorus in the Global Environment; Transfers, Cycles and Management. Scope 54*, John Wiley & Sons, New York, pp. 139–147.
- Lugo A.E., Brown S. and Chapman J. 1988. An analytical review of production rate and stem wood biomass of tropical forest plantations. *Forest Ecol. Manag.* 23: 179–200.
- Malik R. and Sharma S. 1990. Moisture extraction and crop yield as a function of distance in a row of *Eucalyptus tereticornis*. *Agroforest. Syst.* 12: 187–195.
- Mazzarino M.J., Oliva L., Nunez A., Nunez G. and Buffa E. 1991. Nitrogen mineralization and soil fertility in the dry Chaco ecosystem (Argentina). *Soil Sci. Soc. Am. J.* 55: 515–522.
- Mengistu S., Keftasa D. and Yami A. 2002. Productivity of four *Sesbania* species on two soil types in Ethiopia. *Agroforest. Syst.* 54: 235–244.
- Murphy J. and Riley J.P. 1962. A modified single solution method for determination of phosphate in natural waters. *Anal. Chim. Acta* 27: 31–36.
- Ndoye I. and Dreyfus B. 1988. N₂ fixation by *Sesbania rostrata* and *Sesbania sesban* estimated using ¹⁵N and total N different methods. *Soil Biol. Biochem.* 20: 209–213.
- Olsen S.R. and Sommers L.E. 1982. Phosphorus. In: Page A.L. (ed.), *Methods of Soil Analysis, Agron. No. 9, Part 2: Chemical and Microbiological Properties*, 2Am. Soc. Agron., Madison WI., USA, pp. 403–430.
- Palm C.A., Myers R.J.K. and Nandwa S.M. 1997. Combined use of organic and inorganic nutrient sources for soil fertility maintenance and replenishment. In: Buresh R.J., Sanchez P.A. and Calhoun F. (eds), *Replenishing Soil Fertility in Africa. Soil Science Society of America Special Publication 51*, Soil Science Society of America, Madison, WI, USA, pp. 193–218.
- Richards L.A. 1954. *Diagnosis and Improvement of Saline and Alkali Soils. USDA Agric. Handbook 60*, Washington, DC.
- SAS Institute. 1999–2001. SAS/STAT user's guide, release 8.2. SAS Institute Inc., Cary, NC, USA.
- Tedla A., Mamo T., Klajj M.C. and Diedhiou M.L. 1999. Effects of cropping system, seed bed management and fertility interactions on biomass of crops grown on a Vertisol in the central highlands of Ethiopia. *J. Agron. Crop Sci.* 183: 205–211.
- Wood P.J. and Burley J. 1991. A Tree for All Reasons: the Introduction and Evaluation of Multipurpose Trees for Agroforestry. ICRAF, Nairobi, pp. 45–55 and 98–117.